

INTERvention[®]

Farm Animal Care
Disinfectant Cleaner & Deodorizer

Ready to Use Wipes

KEEP OUT OF REACH
OF CHILDREN

FOR BARN
& FARM USE

EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1
EPA Est. No. 075272-WI-001

Reference Sheet

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578
virox.com



This product is a one-step⁸ disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile and stainless steel associated with high touch surfaces, animal equipment, animal crates, automatic feeders[†], buckets, brushes, exterior surfaces of watering appliances, farm vehicles, feed racks[†], foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs and any hard non-porous washable surface where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Claims

*VIRUCIDAL: 1 MINUTE, UNLESS OTHERWISE NOTED

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Avian Reovirus
Bovine Viral Diarrhea Virus
Cytomegalovirus
Duck Hepatitis B virus
Equine Herpes virus type 1
Equine Influenza A Virus (H3N8) (EIV)
Foot and Mouth Disease virus (FMDv) – 3 minutes
Hepatitis B Virus (Duck Hepatitis B virus as the surrogate)
Hepatitis C Virus (Bovine Viral Diarrhea virus as the surrogate)
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B
Human Coronavirus
Influenza A Virus
Influenza B Virus
Norovirus (Feline Calicivirus as the surrogate)
Marek's Disease virus
Newcastle disease virus
Pandemic Influenza A Virus H1N1
Parainfluenza Virus Type 3
Porcine Epidemic Diarrhea virus
Porcine Respiratory & Reproductive Syndrome virus
Porcine Rotavirus
Pseudorabies virus
Respiratory Syncytial Virus (RSV)
Rhinovirus Type-37
Rotavirus (WA strain)

SARS-Related Coronavirus 2 (Human Coronavirus SARS-CoV-2 (the virus that may cause COVID-19⁹))

Vaccinia Virus
Vesicular Stomatitis virus

BACTERICIDAL: 1 MINUTE

Acinetobacter baumannii (Multi-Drug Resistant)
Bordetella bronchiseptica
Bordetella pertussis
Burkholderia cepacia
Enterobacter aerogenes
Enterobacter cloacae (New Delhi metallo-β-lactamase 1 (NDM -1))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Escherichia coli
Escherichia coli O157:H7
Escherichia coli (Extended Spectrum Beta-lactamase (ESBL))
Escherichia coli (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Legionella pneumophila
Listeria monocytogenes
Pasteurella multocida
Pseudomonas aeruginosa
Pseudomonas aeruginosa (Metallo-Beta lactamase (MBL) positive)
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Serratia marcescens
Shigella dysenteriae
Staphylococcus aureus
Staphylococcus aureus Community Acquired Methicillin Resistant Genotype 400 (CA-MRSA 400)
Staphylococcus aureus (Methicillin Resistant (MRSA))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Staphylococcus pseudintermedius
Stenotrophomonas (Pseudomonas) maltophilia
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Streptococcus pyogenes

FUNGICIDAL: 30 SECONDS, UNLESS OTHERWISE NOTED

Candida albicans – 1 minute
Microsporium canis – 1 minute
Trichophyton interdigitale

TUBERCULOCIDAL: 1 MINUTE

Mycobacterium bovis (BCG)

NON-FOOD CONTACT SANITIZER: 10 SECONDS

Acinetobacter baumannii

Campylobacter jejuni

Enterobacter aerogenes

Enterococcus faecalis (Vancomycin Resistant (VRE))

Escherichia coli O157:H7

Listeria monocytogenes

Pseudomonas aeruginosa

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

Mildewcidal Activity – Kills the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity – Controls and prevents the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Malodor Activity – Eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

Re-sealable Soft Pack

1. Remove the seal.
2. Remove one sheet at a time.
3. Close the seal to prevent drying between uses.

Animal Housing Facilities in Veterinary Clinics, Hospitals, Farm Premises including, Equine production Farm, Bovine production Farm.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including pens, stalls, chutes, runs and barns occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
5. All surfaces must remain visibly wet for 1 minute[†].
6. Allow to air dry or rinse with potable water if necessary.
7. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/or Deodorizer

Pull towelette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

For Use as a One-Step Cleaner/Disinfectant

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain visibly wet for 1 minute.[‡]
4. Wipe surface dry or allow to air dry.

***KILLS HIV-1 AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of (inanimate) hard non-porous surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain visibly wet for 1 minute to kill HIV-1, HBV and HCV.

Disposal of Infectious Material: Cleaning materials used that may contain blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Kill Mold and Mildew

1. Pre-clean surfaces.
2. Pull towelette from canister and wipe onto hard, non-porous surfaces until completely wet.
3. Allow the surface to remain visibly wet for 4 minutes.
4. Allow to air dry.

To Control Mold and Mildew

1. Pre-clean surfaces.
2. Pull towelette from canister and wipe onto hard, non-porous surfaces until completely wet.
3. Allow the surface to remain visibly wet for 4 minutes.
4. Allow to air dry.
5. Repeat application weekly or when growth reappears.

For use as a Non-Food Contact Sanitizer

1. Pre-clean visibly soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain visibly wet for 10 seconds.
4. Wipe surface dry or allow to air dry.
5. Change towelette after each use.

Storage And Disposal

PESTICIDE STORAGE: Store this product in a cool dry area, away from direct sunlight and heat. Do not contaminate water, food or feed by storage and disposal.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush. Tightly close lid/flap between uses to retain moisture. Non-refillable container/dispenser.

Do not reuse or refill this container/dispenser. Offer empty container/dispenser for recycling. If recycling is not available, discard container/dispenser in trash.

[†] Rinse with potable water after disinfecting.

[‡] Use a 3min contact time for FMDv.

[§] On hard, non-porous surfaces.

[¶] When used according to disinfection use directions.

^{**} Accelerated Hydrogen Peroxide[®] and Design are registered trademarks of Diversey Inc. | "Accelerated" does not imply increased product efficacy. | For relevant patents, visit: virox.com/patents.